



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

URUGUAY.

Montevideo quarantines against Buenos Ayres and Rosario on account of plague.

The United States consul at Montevideo reports, March 12, that Montevideo has established a quarantine of ten days against Buenos Ayres and Rosario, and five days against the rest of the Argentine ports, as a precaution against plague.

FOREIGN STATISTICAL REPORTS.

AUSTRIA—*Trieste*.—Year ended December 31, 1899. Estimated population, 166,745. Total number of deaths, 4,605, including diphtheria, 109; enteric fever, 42; scarlet fever, 165, and 769 from tuberculosis.

ECUADOR—*Guayaquil*.—Ten days ended April 1, 1900. Estimated population, 55,000. Total number of deaths, 146. No contagious diseases reported.

FRANCE—*Nice*.—Four weeks ended April 10, 1900. Estimated population, 108,227. Total number of deaths, 171, including diphtheria, 1; enteric fever, 1, and 4 from smallpox.

Roubaix.—Month of March, 1900. Estimated population, 127,327. Total number of deaths, 250, including diphtheria, 4, and 2 from measles.

GERMANY—*Dresden*.—Month of February, 1900. Estimated population, 402,400. Total number of deaths, 497, including diphtheria, 4; measles, 1; whooping cough, 5, and 81 from phthisis pulmonalis.

Hanover.—Month of January, 1900. Estimated population, 247,140. Total number of deaths, 312, including diphtheria, 3; enteric fever, 1; scarlet fever, 1, and 41 from phthisis pulmonalis.

Month of February, 1900. Estimated population, 247,593. Total number of deaths, 404, including diphtheria, 3; enteric fever, 1; scarlet fever, 3, and 49 from phthisis pulmonalis.

Weimar.—Month of March, 1900. Estimated population, 29,633. Total number of deaths, 48. No contagious diseases reported.

GREAT BRITAIN—*England and Wales*.—The deaths registered in 33 great towns in England and Wales during the week ended March 31, 1900, correspond to an annual rate of 21.4 a thousand of the aggregate population, which is estimated at 11,610,296. The highest rate was recorded in Wolverhampton, viz, 33.8, and the lowest in Croydon, viz, 11.9.

London.—One thousand seven hundred and eighty-five deaths were registered during the week, including measles, 60; scarlet fever, 7; diphtheria, 31; whooping cough, 45; enteric fever, 7; and diarrhea and dysentery, 18. The deaths from all causes correspond to an annual rate of 20.3 a thousand. In Greater London 2,381 deaths were registered, corresponding to an annual rate of 18.7 a thousand of the population. In the "outer ring" the deaths included 18 from diphtheria, 9 from measles, 3 from scarlet fever, and 15 from whooping cough.